

Bay of Bengal Programme

Development of Small-Scale Fisheries

GROWTH AND MORTALITY OF THE MALAYSIAN
COCKLE (ANADARA GRANOSA L.) UNDER
COMMERCIAL CULTURE:
ANALYSIS THROUGH LENGTH-FREQUENCY DATA

BOBP/WP/47



SWEDISH INTERNATIONAL DEVELOPMENT AUTHORITY



FOOD AND AGRICULTURE ORGANISATION OF THE UNITED NATIONS

BAY OF BENGAL PROGRAMME
Development of Small-Scale Fisheries

BOBP/WP/47
GCP1 RAS/040/SWE

GROWTH AND MORTALITY OF THE MALAYSIAN
COCKLE (*ANADARA GRANOSA* L.) UNDER
COMMERCIAL CULTURE:
ANALYSIS THROUGH LENGTH-FREQUENCY DATA

BOBP/WP/47

by Ng Fong Oon
Fisheries Research Institute
Glugor, Penang, Malaysia

Executing Agency:
Food and Agriculture Organization
of the United Nations

Funding Agency:
Swedish International
Development Authority

Development of Small-Scale Fisheries in the Bay of Bengal. Madras, India, July 1986
Mailing Address: Post Bag. 1054, Madras 600018, India
Street Address: 91 St. Mary's Road, Abhirampuram, Madras 600018, India
Cables: FOODAGRI Telex: MS-311 FISH Phones: 71294, 71296, 71587, 77760

This paper analyses length frequency data, by several methods, of the Malaysian cockle (*Anadara granosa* L). The data were collected monthly from five different plots under commercial culture during a period of 12-17 months. Parameters of the von Bertalanffy growth formula (VBGF) were derived for each of the five plots, along with estimates of related parameters (mortality, mean length of first capture, etc.). "Yield per recruit" analyses suggest that the present legal size for the five culture plots is well above the maximum yield per recruit. The paper discusses the limitations of the methodology and data used. Suggestions for further studies are also made.

The cockle samples were collected and measured by staff of the cockle team, headed by the author, of the Glugor Fisheries Research Station (GFRI), Penang, Malaysia. The analyses were made by the author on a fellowship visit to the International Center for Living Aquatic Resources Management (ICLARM), Manila, in October 1985.

The author wishes to thank the GFRI's Director of Research, Mr. Mohd Shaari bin Sam Abdul Latiff, and Mr. Ong Kah Sin, Head of the Aquaculture Section, for their encouragement and suggestions on the project; Dr. Daniel Pauly (ICLARM), for his help with the analyses; Dr. J. Saeger and Mr. Gayanilo (GTZ) for allowing him to use their revised version of the ELEFAN programs and their computer facilities; Ms. Faazaz bte Latiff, Mr. Kamal Zaman bin Mohamad and Ms. Devaki Nair for collecting and compiling the length frequency data at Penang and Set an gor.

The work described in this paper is one component of a programme for the Development and Management of Cockle Culture in Malaysia, supported by the small-scale fisheries project of the Bay of Bengal Programme (BOBP). The programme's first phase, undertaken during 1985, consisted primarily of biological studies. Other components were experiments with induced spawning, determination of maturity and spawning pattern by means of condition indices, monitoring of culture plots, and examination of cockle shells to determine age and growth pattern. The next phase will consist of economic and socio-economic studies.

The small-scale fisheries project of the Bay of Bengal Programme began in 1979 and covers five countries bordering the Bay of Bengal – Bangladesh, India, Malaysia, Sri Lanka and Thailand. Funded by SIDA (Swedish International Development Authority) and executed by the FAO (Food and Agriculture Organization of the United Nations), the project seeks to develop, demonstrate and promote appropriate technologies and methodologies to improve the conditions of small-scale fisherfolk in member countries.

This document is a working paper and has not been officially cleared by the Government concerned or by the FAO.

CONTENTS

| | <i>Page</i> |
|----------------------------------|-------------|
| 1. Introduction | 1 |
| 2. Location of the culture plots | 1 |
| 3. Material and method | 2 |
| 4. Results | 4 |
| 5. Discussion | 5 |
| 6. References | 6 |

Figures

| | |
|--|----|
| 1. Map of peninsular Malaysia showing locations of culture plots A to E | 8 |
| 2. Estimation of L_{∞} and K using ELEFAN I, Plot A | 9 |
| 3. Catch curve and selection pattern for <i>A. granosa</i> , Plot A | 10 |
| 4. Estimation of L_{∞} and Z/K using the method of Wetherall <i>et al.</i> , Plots A to E | 11 |
| 5. Yield-per-recruit analyses of <i>A. granosa</i> , Plots A to E | 12 |

Appendix

| | |
|--|----|
| 1. Length frequency data of <i>Anadara granosa</i> from five culture plots | 13 |
|--|----|

| | |
|--|----|
| <i>Publications of the Bay of Bengal Programme</i> | 18 |
|--|----|

1. INTRODUCTION

Cockle culture plays a major economic role in Malaysia: it accounts for about 11% of the country's total fisheries production. In 1984 about 65,000 tonnes of cockles were produced (Anon. 1985). It is by far the most important aquaculture industry in Malaysia. Currently about 4,000 to 5,000 hectares of mudflats along the west coast of peninsular Malaysia are utilized for the culture of this important bivalve.

There have been, however, great fluctuations in the production of cockles and in prices of spats in the recent past. Production reached an all-time high of 120,000 tonnes in 1980; it was only about 40,000 tonnes in 1983. The prices of spats more than doubled from 1979 to 1980 and remained high for three years. The increase in price, an indication of shortage of spats, prompted the Government to intervene to stabilize the industry and prevent overexploitation of the cockle population. The existing legislation on minimum size of harvested cockles was strictly enforced, and a ban imposed on the export of spats (mainly to Thailand). The minimum harvesting size of 31.8 mm has met with much opposition from the cockle farmers since they consider it too large for viable culture operations.

The Government has, therefore, relaxed the enforcement of the legislation and launched a programme of biological and economic studies to obtain more information as a basis for perhaps more appropriate management measures. This programme is conducted with support from the BOBP.

The general biology and culture aspects of the Malaysian cockle (*Anadara granosa*) have been well described by Pathansali and Soong (1958), Pathansali (1963, 1966), Broom (1980, 1982a, 1982b, 1982c, 1982d, 1983a, 1983b, 1983c), Ng (1982, 1984) and Wong and Lim (1985). But information on growth and mortality in general and under different culture conditions is still very scanty. The study presented in this paper addresses itself to this particular aspect of cockle biology. Five culture plots in important cockle producing areas were selected to study growth and mortality parameters by using length-frequency data.

While several methods are available to the fishery biologist for investigating growth and mortality patterns, the computer-based ELEFAN method was selected, because it obviates the need for precise and reliable production estimates and age determinations. The methodology for age determination of the cockles had not been devised at the start of this project, nor had any adaptations been made from that used for other shellfishes. Moreover, age determinations require much more trained manpower than does the establishment of length-frequency sampling.

2. LOCATION OF THE CULTURE PLOTS

Five commercial culture plots were selected for the study. Three plots were selected in the state of Perak, and one each in the states of Penang and Selangor. Their geographic locations are shown in Figure 1.

Culture plot "A" is located outside and south of the river mouth, Kuala Juru, Penang. The surface area of the culture plot is about 20 ha and it is located between Mean High Water of Neap (M.H.W.N.) and Mean Low Water of Neap tides (M.L.W.N.).

"B" of about 40 ha, is located about one km off Pulau Sangga Besar, Perak and the culture site is between M.H.W.N. and M.L.W.N.

"C", is located in Perak just inside the mouth of the river Kuala Larut. It has an area of about 20 ha and is located between M.L.W.N. and Mean Low Water Sprint tide (M.L.W.S.).

“D”, is also located in the river Kuala Larut about 0.5 km inside the river mouth between M.H.W.N. and M.L.W.N. The size is about 40 ha.

“E”, is located about one kilometre outside the river Sungei Besar, Selangor, between M.H.W.N. and M.L.W.N.

Except with regard to salinity and suspended solids, the ecological characteristics of the five culture plots did not differ significantly. In the case of salinity, the culture plots at A, C and D seemed to be subjected to a wider range of fluctuation than B and E. Total suspended solids values at C and D were relatively high during the month of May. Details concerning ecological characteristics are given in the table below.

Ecological characteristics of culture plots A—E

| Parameter | Culture Plot | | | | |
|--------------------------------|--------------|-------------------------------|-------------------------|-------------------------|-------------------------|
| | A | B | C | D | E |
| Salinity (%) | 16—31 | 22—29 | 14—27 | 12—25 | 22—30 |
| Water pH | 7.9—8.5 | 7.6—8.3 | 7.1—8.0 | 7.1—8.0 | 7.8—8.4 |
| Dissolved O ₂ (ppm) | 5.6—9.1 | 6.2—11.0 | — | — | — |
| Mud organic content (%) | — | 9.1—15.2 | 6.2—19.1 | 6.2—19.1 | — |
| Suspended solid (ppm) | — | 10—160 | 10—400 | 10—400 | — |
| Soil type | sandy | sandy loam to sandy clay loam | sandy loam to clay loam | sandy loam to clay loam | sandy loam to clay loam |

3. MATERIAL AND METHOD

Monthly random samples of about 500 cockles, mud and water were taken from each of the five culture plots. The sampling was conducted from May 1984 till September 1985 and the duration for each plot was as follows:

| | |
|------|----------------------------|
| A | June 1984—September 1985 |
| B, C | September 1984—August 1985 |
| D | May 1984—September 1985 |
| E | July 1984—September 1985 |

Length and weight measurements were made using a vernier caliper (to the nearest 0.1 mm) and a digital balance (0.1 g), respectively. Random samples were taken instead of taking samples from the same predetermined sites within each plot in order not to reduce the density of any one area within the culture plot. Generally, a minimum of about 3 sub samples (150 cockles) were taken from each culture plot. The length-frequency data are given in Appendix 1.

The length-frequency data were analyzed using the ELEFAN I and II programs (Pauly 1984, Pauly and David 1981, Pauly, *etal.* 1984) and the revised version of the ELEFAN I and II programs by Saeger and Gayanilo (1985).

Also, the method of Wetherall *et al.* (1986) was used to estimate L_{∞} and Z/K from the length-frequency data.

A comparison of the growth performance of *A. granosa* in the five culture plots was carried out using the parameter ϕ as defined in Pauly and Munro (1984), i.e.:

$$\phi = \log_{10} K + 2 \log L_{\infty} \dots \dots (1)$$

where K and L_{∞} are parameters of the von Bertalanffy growth formula, (VBGF) stands for length and \log stands for decimal logarithm.

Pending a detailed analysis of the available length-weight data, a preliminary length-weight relationship of the form

$$W = a.L^3 \dots \dots (2)$$

was derived, using 10 pairs of length and weight measurements from plot D, pertaining to the 10 largest specimens measured at that plot.

Equation (2) was then used to convert the available estimate of L_{∞} into estimates of W_{∞} .

Values of the parameter t_0 of the VBGF were obtained finally by solving it, for each plot, for an age of 4 months, i.e., 1/3 year at 6.4 mm, the legal size limit for seeds (see Broom 1983a).

Thus, for each plot, estimates of the 3 parameters of the VBGF for growth:

$$W_{\infty} = W_{\infty} \left(1 - e^{-k(t-t_0)} \right) \dots \dots (3)$$

are available, such that standard yield per recruit analyses (Beverton and Holt 1957, Pauly 1984) can be performed. Since values of natural mortality (M) in *Anadara granosa* are not available, the yield per recruit analyses was performed for a range of M values ($M=1/2F$, $M=F$, $M=2F$), with the sum of fishing (F) and natural mortality being constrained by the total mortality ($Z=F+M$) observed at each site.

4. RESULTS

The growth, mortality and related parameters of the five culture plots, (L_{∞} , K , Z , L_{50} , L' and ϕ') obtained from the detailed analysis of the length-frequency data by ELEFAN (I and II) are tabulated below:

Estimates of growth and mortality parameters of *Anadara granosa* from five culture plots using elefan I and II.

| | Plot | | | | |
|---------------------------|-------|-------------|-------------|--------|-------|
| | A | B | C | D | E |
| L_{∞} (mm) | 45.0 | 37.4 | 40.5 | 34.2 | 41.4 |
| W_{∞} (g) | 27.3 | 15.7 | 19.9 | 12.0 | 21.3 |
| K (1/yr) | 0.55 | 0.87 | 0.79 | 0.60 | 0.78 |
| t_0 (yr) | 0.054 | 0.118 | 0.116 | -0.112 | 0.118 |
| Z (1/yr) | 3.24 | 2.93 | 4.12 | 4.04 | 3.66 |
| L_{min} (mm) | 7.0 | 10.5 | 16.5 | 5.5 | 7.0 |
| L_{50} (mm) | 25.6 | 21.5 | 26.3 | 19.4 | 27.6 |
| L' | 26.0 | 22.0 | 29.0 | 20.0 | 28.0 |
| ϕ' | 3.047 | 3.085 | 2.846 | 2.846 | 3.126 |
| Legal age at harvest (yr) | 2.28 | 2.30 | 2.06 | 4.42 | 1.99 |

- W_{∞} — Computed by using preliminary length-weight relationship based on data from plot D i.e. $W=0.0003 L^3$.
- t_0 — Computed by solving the VBGF for an age of 4 months at 6.4 mm.
- L_{min} — Minimum size in samples.
- L_{50} — Length at which probability of capture is 50 per cent (using a "Kor", i.e. the hand-held dredge used to collect samples).
- Legal age at harvest — Converted from legal length of 31.8 mm and growth parameters L_{∞} , K and t_0 .

The value of ϕ' showed that the cockles in culture plots A and B are similar in their growth performance; likewise for the cockles in plots C and D. These values of ϕ' imply that the cockles cultured at plots A and B grow better than cockles in plots C and D. Figure 2 shows an example (for Plot A) of the restructured length-frequency data and the optimum growth curves fitted by ELEFAN I. A good fit of the growth curve to the succession of runs of positive values (peaks) of the length-frequency data was attained for all five culture plots.

In each culture plot there is only one year class with the exception of some odd older animals remaining from previous culture cycles. The presence of two distinct year classes is thus an artifact due to the presentation of the data; the sampling lasted more than a year resulting in an overlapping of data in plots A, D and E.

The catch curve and selection pattern as calculated by ELEFAN II for data from Plot A is shown as an example in Fig. 3.

The estimates of L_{∞} and Z/K using the method of Wetherall *et al.* (1986) for plots A to E are given in Fig. 4.

The L_{∞} values obtained by the two methods compared favourably and appeared almost identical for culture plots A, C, and D. Details are given in the table below.

Comparison of parameter estimates using ELEFAN (I&II) and method of Wetherall *et al.*, (1986).

| Parameter Method | L_{∞} | | Z/K | |
|---------------------|--------------|-----------|---------|-----------|
| | ELEFAN | WETHERALL | ELEFAN* | WETHERALL |
| A | 45.0 | 46.4 | 5.891 | 3.444 |
| B | 37.4 | 34.7 | 3.368 | 2.531 |
| C | 40.5 | 38.4 | 5.215 | 4.128 |
| D | 34.2 | 34.0 | 6.773 | 4.914 |
| E | 41.4 | 42.1 | 4.692 | 4.462 |

* From catch curve estimates (Fig. 3).

The yield per recruit analyses (Fig. 5) all suggest that yield per recruit is maximized, for each of the five culture plots, by a minimum age of harvest of about one year, well below the age corresponding to the legal harvesting size.

5, DISCUSSION

The growth and mortality parameters for cockles in the five culture plots clearly indicate differences between plots. It is difficult, at this stage of the study, to interpret these differences. Though they were expected because of known dissimilarities in ecological features of the five culture plots, these differences could also be due to other factors that were not recorded, such as density differences or a size distribution that is related to water depth.

The use of the parameter ϕ' furthermore confirmed that the cockles in culture plots A, B and E were similar in growth pattern; a similar growth pattern was observed for culture plots C and D. The values of ϕ' showed that the cockles cultured at plot A, B and E grew better than those in plots C and D.

Comparison of the growth parameter L_{∞} estimated by ELEFAN I and by the method of Wetherall *et al.* (1986) shows values which are similar, especially in culture plots A, C, D and E. The point to note, however, is that the estimates of Z/K obtained by both methods show marked differences. These differences are not surprising in view of the fact that in both cases, the final values obtained are largely influenced by the points included in the computations. Despite these discrepancies, the estimates of L_{∞} and Z/K obtained by both methods are consistent as a whole.

Another observation worthy of note is that the growth curve estimated by ELEFAN I seems not to have been affected by the constant shifting of cockles from one end of the culture plot to the other. The yield per recruit analyses carried out on each of the five culture plots all suggest that the yield per recruit is maximized by a minimum age at harvest of about one year, well below the age corresponding to the present legal harvesting size of 31.8 mm.

One could question the validity of the values of parameter W_{∞} for the five culture plots, since these values were computed on the basis of length-weight relationships from culture plot D only.

However, yield per recruit estimates are directly proportional to W_a ; hence using different values of this parameter would only change the ordinate scale of the graphs in Fig. 5, but not the overall shape of the yield curves, which would still suggest that optimum harvesting size is obtained after one year.

Figure 5 shows that yield per recruit is independent of the fishing mortality, over the range of mortalities considered here. In other words, the yield per recruit analysis clearly indicates that it is indeed uneconomic to grow cockles for more than one year because of diminishing returns. The evidence from the yield per recruit analysis suggests that the present regulation on size limit on harvesting cockles should be reappraised if returns *on a yield per recruit basis* are to be *maximized*. While accepting the fact that the regulation on size limit is based on sexual maturity as well as the possibility of the cockles spawning at least once prior to their harvest, there is the need to consider the economic factor as well. Moreover, there is no evidence to support the assumption that the yearly production of seeds is directly related to the size of the *cultured* broodstock. Most culture practices today depend largely on natural seed supplies. The fluctuating pattern of seed supplies is universal and it depends largely on the environmental conditions which prevail in each country and area (Broom 1985).

The assumption that retaining a substantial broodstock in the culture areas would increase the yield per recruit could be counter-productive as well. Cockles are filter-feeders and we can therefore assume that a significant proportion of newly released gametes are filtered out by the broodstock when the cockles are kept in a crowded situation (i.e. before they reach the minimum legal size of harvest).

While a substantial amount of information has been gained by the analyses of data using the ELEFAN programmes, particularly on growth mortality and yield per recruit, further studies on possible causes of recruitment variability would go a long way in providing the answer to the question related to seed supply and production. These studies, together with other on-going studies on maturation and spawning, and possibly also a study on genetic variability, will further provide the necessary information on which future management decisions could be based.

REFERENCES

- Anon. 1985. Annual States Fisheries Statistics. 1984 Kuala Lumpur, Malaysia, Department of Fisheries, Mm. of Agriculture.
- Beverton, R.J.H. and S.J. Holt. 1957. On the dynamics of exploited fish populations. Fish. Invest. Ser. II. No. 19. 533 p.
- Broom, M.J. 1980.** Community and production ecology of *Anadara granosa* (L) with particular reference to its gastropredators. Kuala Lumpur, University of Malaya. 349 p. Ph.D. thesis.
- Broom, M.J. 1982a.** Structure and seasonality in a Malaysian mudflat community. *Estuarine, Coastal and Shelf Science*. 15: 135-150.
- Broom, M.J. 1982b.** Analysis of the growth of *Anadara granosa* (L) (Bivalvia: Arcidae) in natural artificially seeded and experimental populations, *Marine Ecology Progress Series* 9: 69-79.
- Broom, M.J. 1982c. Size selection, consumption rates and growth of the gastropods *Nat/ca maculosa* Lamarck and *Thais carinifera* (Lamarck) preying on the bivalve *Anadara granosa* (L). *Journal of Experimental Marine Biology and Ecology*. 56: 213-233.
- Broom, M.J. 1982d.** The management of *Anadara granosa* (L) as a natural resource. *Resource Management Optimization* 2: 1-23.
- Broom, M.J. 1983a. Mortality and production in natural artificially seeded and experimental populations of *Anadara granosa* (L) (Bivalvia: Arcidae). *Oecologia* (Berlin) 58: 389-397.

- Broom, M.J. 1983b.** A preliminary assessment of prey species preference by the tropical marine gastropods (*Natica maculosa* Lamarck and *Thais carinifera* (Lamarck)). *Journal of Molluscan Studies*. 49: 43-52.
- Broom, M.J. 1983c. Gonad development and spawning in *Anadara granosa* (L) (Bivalvia: Arcidae) *Aquaculture* 30: 211-219.
- Broom, M.J.** 1985. Biology and culture of marine bivalves molluscs of the genus *Anadara*. ICLARM Stud. Rev. (in press).
- Ng, F.O.** 1984. Cockle culture. Bangkok. Southeast Asian Fisheries Information Service. Extension Manual Series No. 13: 22 p.
- Ng, F.O., **J. Pang** and **T.P. Tang.** 1982. Country report: Malaysia p.47-52 In: (eds) by F. Brian Davy and M. Graham. Bivalve Culture in Asia and the Pacific at Singapore. International Development Research Centre. IDRC-200e. 90 p.
- Pauly, D. 1984. Fish population dynamics in tropical waters: a manual for use with programmable calculators. Manila ICLARM Stud. Rev. 8: 325 p.
- Pauly, D. and N. **David.** 1981. ELEFAN I, a BASIC programme for the objective extraction of growth parameters from length frequency data. *Meeresforschung* 28(4): 205-211.
- Pauly, D., J. Inglos and R. Neal.** 1984. Application to shrimp stocks of objective methods for the estimation of growth, mortality and recruitment-related parameters from length data. p. 220-234. In Penaeid shrimps: their biology and management ed. by J.A. Gulland and B.J. Rothschild. 220-234 Farnham, England Fishing News Books, 308 p.
- Pauly, D. and J.L. Munro.** 1984. Once more on the comparison of growth in fish and invertebrates. *Fishbyte* 2(1): 21.
- Pathansali, D. 1963.** On the effect of salinity changes on the activity of the cockle, *Anadara granosa* (L). *Malaysian Agriculture Journal* 44: 18-25.
- Pathansali, D. 1966. Notes on the biology of the cockle *Anadara granosa* (L). Proceedings of Indo-Pacific Fishery Council 11: 84-98.
- Pathansali, D. and M.K. Soong. 1958. Some aspects of cockle (*Anadara granosa* L.) culture in Malaysia. Proceedings of Indo-Pacific Fishery Council. 8: 26-31.
- Saeger, J. and F.C. Gayanilo.** 1985. ELEFAN I and II live on screen, *Fishbyte* 3(2): 13-14.
- Wetherall, J.A., J.J. Polovina and S. Ralston.** 1986. Estimating growth and mortality in steady state fish stocks from length-frequency data. In: Length-Based Methods in Fishery Research ed. by D. Pauly and G.R. Morgan. ICLARM Conference Proceedings (in press).
- Wong, T.M. and **Lim. T.G.** 1985. Induced spawning of *Anadara granosa* in Malaysia. *ICLARM Newsletter* B(4) (in press).

Figure 1
MAP OF PENINSULAR MALAYSIA
SHOWING LOCATION OF CULTURE PLOTS A TO E.




Figure 2

ESTIMATION OF L_a AND K USING ELEFAN I, PLOT A

Note good fit of the growth curve to the succession of runs of positive values (i.e., peaks) of the length-frequency data in Appendix 1, as restructured by Elefan 1.




Figure 3

CATCH CURVE AND SELECTION PATTERN FOR *A. Granosa*, PLOT A.

The slope of the catch provides an estimate of total mortality (z), while the ascending, left side of the curve leads to the selection pattern.

Ancdaro granosa
K JuruPenang
 $L_{\infty} = 45$, $K = .55$




Figure 4
ESTIMATION OF L_{∞} AND Z/K USING THE METHOD OF WETHERALL *ET AL.* (1986) PLOTS A TO E.
Based on length-frequency data in Appendix 1 each combined into a single, mean annual sample.





Figure 5

YIELD-PER-RECRUIT ANALYSES OF *A. Granosa*, PLOTS A TO E

Note that all five plots suggest that the present minimum legal size (i.e., age) at harvest is too high (particularly in cases of s/o/w growth such as in Plot D), and that this result is independent of F and M over the range considered here, and of specific values of W_a .



Appendix 1

Length-frequency data of *Anadara granasa* from five culture plots Plot A. K. Juru, Penang.

| Size (mm) | Sampling date | | | | | | | | | | | |
|--------------|---------------|------|------|------|------|------|------|------|------|-------|-------|-------|
| | 1/15 | 2/15 | 3/15 | 4/15 | 5/15 | 6/15 | 7/15 | 8/15 | 9/15 | 10/15 | 11/15 | 12/15 |
| 7.0 | | | | | | 120 | 11 | 1 | | | | |
| 9.0 | | | | | | 734 | 116 | 41 | 1 | 2 | | |
| 11.0 | | | | | | 580 | 257 | 120 | 11 | 8 | | |
| 13.0 | 8 | 1 | | | | 70 | 180 | 135 | 65 | 18 | 6 | 1 |
| 15.0 | 89 | 25 | | 2 | 14 | 1 | 78 | 68 | 73 | 24 | 44 | 25 |
| 17.0 | 348 | 122 | 26 | 16 | 38 | 0 | 53 | 21 | 45 | 89 | 200 | 75 |
| 19.0 | 449 | 89 | 140 | 106 | 98 | 1 | 37 | 26 | 9 | 76 | 138 | 131 |
| 21.0 | 219 | 38 | 237 | 234 | 100 | 10 | 36 | 54 | 34 | 31 | 28 | 104 |
| 23.0 | 34 | 34 | 203 | 323 | 107 | 51 | 66 | 72 | 68 | 7 | 1 | 34 |
| 25.0 | 2 | 47 | 84 | 237 | 111 | 127 | 124 | 172 | 108 | 20 | 5 | 29 |
| 27.0 | | | 30 | 78 | 96 | 169 | 158 | 219 | 127 | 19 | 27 | 52 |
| 29.0 | | | 16 | 15 | 63 | 127 | 128 | 162 | 102 | 17 | 57 | 64 |
| 31.0 | | | 4 | 3 | 22 | 37 | 99 | 43 | 55 | 2 | 58 | 37 |
| 33.0 | | | | | 9 | 21 | 56 | 13 | 23 | 1 | 23 | 9 |
| 35.0 | | | | | 1 | | 20 | 3 | 6 | | 5 | 9 |
| 37.0 | | | | | | | 1 | 5 | 1 | | 1 | 4 |
| 39.0 | | | | | | | | 1 | | | 3 | 4 |
| 41.0 | | | | | | | | 1 | | | 2 | |
| 43.0 | | | | | | | | | | | 1 | |

Appendix 1 (Continued)

Length-frequency data of *Anadara granosa* from five culture plots.

Plot B: B1, Pulau Sangga, Perak

| Size (mm) | Sampling date | | | | | | | | | | | |
|--------------|---------------|------|------|------|------|------|------|------|------|-------|-----------|-----------|
| | 1/30 | 2/27 | 3/29 | 4/25 | 5/15 | 6/15 | 7/31 | 8/28 | 9/15 | 10/31 | 11/28 | 12/28 |
| 10.5 | | | | | | | | | 2 | | | |
| 11.5 | | | | | | | | | 11 | 3 | | |
| 12.5 | | | | | | | | | 43 | 9 | | |
| 13.5 | | | | | | | | | 96 | 16 | | |
| 14.5 | | | | | | | | | 118 | 50 | 2 | |
| 15.5 | 5 | | | | | | | | 114 | 108 | 19 | 2 |
| 16.5 | 13 | 2 | | | | | | | 80 | 156 | 33 | 10 |
| 17.5 | 20 | 5 | 1 | 2 | | | | | 28 | 200 | 36 | 7 |
| 18.5 | 57 | 8 | 5 | 0 | | | | | 5 | 186 | 102 | 11 |
| 19.5 | 81 | 32 | 18 | 4 | | 2 | | | 4 | 102 | 137 | 43 |
| 20.5 | 115 | 70 | 53 | 18 | 4 | 7 | 1 | | | 41 | 162 | 59 |
| 21.5 | 97 | 96 | 80 | 37 | 2 | 24 | 6 | 1 | | 8 | 104 | 67 |
| 22.5 | 71 | 99 | 94 | 80 | 17 | 43 | 10 | 0 | | | 49 | 68 |
| 23.5 | 35 | 64 | 92 | 125 | 42 | 77 | 12 | 1 | | | 22 | 58 |
| 24.5 | 7 | 31 | 54 | 113 | 77 | 111 | 18 | 12 | | | 4 | 39 |
| 25.5 | | 12 | 29 | 77 | 97 | 88 | 31 | 30 | | | | 15 |
| 26.5 | | 1 | 3 | 33 | 94 | 68 | 43 | 43 | | | | |
| 27.5 | | | | 15 | 62 | 26 | 84 | 57 | | | | |
| 28.5 | | | | 2 | 50 | 17 | 88 | 60 | | | | |
| 29.5 | | | | | 14 | 9 | 69 | 61 | | | | |
| 30.5 | | | | | 6 | 6 | 59 | 59 | | | | |
| 31.5 | | | | | 1 | 3 | 20 | 39 | | | | |
| 32.5 | | | | | | 2 | 13 | 23 | | | | |
| 33.5 | | | | | | 1 | 4 | 8 | | | | |

Appendix 1 (Continued)

Length-frequency data of *Anadara granosa* from the five culture plots.

Plot C: A12 K, Sepetang, Perak

| Size (mm) | Sampling data | | | | | | | | | | | |
|--------------|---------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------|----------|----------|-----------|
| | 1/29 | 2/26 | 3/28 | 4/24 | 5/30 | 6/15 | 7/30 | 8/28 | 9/15 | 10/30 | 11/27 | 12/27 |
| 16.5 | | 1 | | | | | | | | | | 16 |
| 17.5 | 1 | 0 | | | | | | | 5 | | | 31 |
| 18.5 | 2 | 2 | | | | | | | 18 | 2 | 2 | 40 |
| 19.5 | 12 | 4 | | 3 | | | | | 25 | 6 | 5 | 55 |
| 20.5 | 13 | 11 | 1 | 7 | | | | 2 | 42 | 12 | 9 | 75 |
| 21.5 | 49 | 23 | 10 | 12 | | 3 | 1 | 2 | 24 | 36 | 37 | 96 |
| 22.5 | 80 | 38 | 10 | 21 | 1 | 0 | 0 | 0 | 16 | 60 | 44 | 65 |
| 23.5 | 93 | 64 | 22 | 42 | 13 | 1 | 0 | 3 | 6 | 81 | 87 | 40 |
| 24.4 | 88 | 85 | 32 | 77 | 19 | 1 | 2 | 6 | 0 | 77 | 78 | 10 |
| 25.5 | 63 | 96 | 33 | 106 | 44 | 19 | 5 | 4 | 1 | 44 | 53 | 8 |
| 26.5 | 24 | 76 | 28 | 91 | 54 | 28 | 19 | 15 | | 17 | 19 | 1 |
| 27.5 | 14 | 43 | 12 | 55 | 53 | 54 | 35 | 58 | | 4 | 5 | 1 |
| 28.5 | | 22 | 6 | 29 | 82 | 72 | 50 | 64 | | | 1 | 1 |
| 29.5 | | 3 | 6 | 22 | 62 | 66 | 64 | 50 | | | | |
| 30.5 | | | | 8 | 56 | 51 | 48 | 52 | | | | |
| 31.5 | | | | 3 | 28 | 32 | 40 | 29 | | | | |
| 32.5 | | | | 2 | 12 | 18 | 26 | 19 | | | | |
| 33.5 | | | | | 14 | 3 | 14 | 8 | | | | |
| 34.5 | | | | | | 4 | 7 | 5 | | | | |
| 35.5 | | | | | | 1 | 7 | 2 | | | | |
| 365 | | | | | | | 1 | 1 | | | | |

Appendix 1 (Continued)

Length-frequency data of *Anadara granosa* from five culture plots.

Plot D: A 15 K, Sepetang, Perak

| Size (mm) | Sampling date | | | | | | | | | | | |
|--------------|---------------|-----------|----------|----------|------------|-----------|-----------|-----------|------------|-----------|-----------|-----------|
| | 1/29 | 2/26 | 3/28 | 4/24 | 5/30 | 6/15 | 7/28 | 8/28 | 9/28 | 10/30 | 11/27 | 12/27 |
| 5.5 | | | | | 6 | 5 | 4 | | | | | |
| 6.5 | | | | | 44 | 23 | 29 | 2 | | | | |
| 7.5 | | | | | 182 | 68 | 74 | 13 | | | | |
| 8.5 | | | | | 321 | 136 | 152 | 75 | | | | |
| 9.5 | | | | | 363 | 179 | 211 | 165 | 10 | | | |
| 10.5 | | | | | 172 | 155 | 195 | 191 | 16 | | | |
| 11.5 | 4 | 2 | | | 59 | 75 | 137 | 162 | 65 | 2 | 2 | |
| 12.5 | 7 | 4 | | | 18 | 38 | 79 | 90 | 116 | 12 | 18 | |
| 13.5 | 17 | 14 | | 1 | 3 | 11 | 30 | 59 | 170 | 29 | 41 | 2 |
| 14.5 | 44 | 47 | 6 | 8 | 3 | 3 | 12 | 24 | 159 | 48 | 79 | 14 |
| 15.5 | 86 | 69 | 15 | 22 | 7 | 2 | 4 | 10 | 115 | 103 | 94 | 63 |
| 16.6 | 113 | 114 | 57 | 28 | 20 | 7 | 15 | 2 | 75 | 103 | 96 | 95 |
| 17.5 | 78 | 80 | 92 | 75 | 35 | 19 | 22 | 4 | 35 | 113 | 76 | 137 |
| 18.5 | 76 | 64 | 114 | 95 | 76 | 55 | 46 | 22 | 27 | 77 | 55 | 91 |
| 19.5 | 31 | 43 | 87 | 102 | 93 | 80 | 78 | 37 | 21 | 37 | 37 | 60 |
| 20.5 | 28 | 29 | 65 | 76 | 113 | 102 | 84 | 62 | 42 | 10 | 5 | 27 |
| 21.5 | 5 | 10 | 38 | 49 | 65 | 65 | 87 | 100 | 31 | 4 | 1 | 13 |
| 22.5 | 1 | 2 | 21 | 25 | 43 | 71 | 77 | 102 | 81 | | | 5 |
| 23.5 | | | 7 | 19 | 34 | 44 | 76 | 66 | | | | 2 |
| 24.5 | | | 4 | 4 | 10 | 11 | 17 | 64 | 62 | | | |
| 25.5 | | | | 5 | 2 | 12 | 0 | 23 | 41 | | | |
| 26.5 | | | | | | 5 | 1 | 8 | 24 | | | |
| 27.5 | | | | | | | 3 | 2 | 16 | | | |
| 28.5 | | | | | | | 1 | | 6 | | | |
| 29.5 | | | | | | | | | 2 | | | |
| 30.5 | | | | | | | | | 3 | | | |

Appendix 1 (Continued)

Length-frequency data of *Anadara granosa* from five culture plots.

Plot E: S. Besar, Se/angor

| Size (mm) | Sampling date | | | | | | | | | | | |
|--------------|---------------|-----|-----|-----|-----|------|------|------|------|------|------|-------|
| | 1/8 | 2/6 | 3/6 | 4/9 | 5/8 | 6/12 | 7/19 | 8/14 | 9/15 | 10/8 | 11/8 | 12/11 |
| 7.0 | | | | | | | 1 | 2 | 4 | | | |
| 9.0 | | | | | | | 39 | 11 | 19 | 17 | 1 | |
| 11.0 | | | | | | | 106 | 38 | 47 | 51 | 10 | |
| 13.0 | | | 1 | | | | 129 | 58 | 89 | 54 | 26 | 1 |
| 15.0 | 15 | | | | 2 | | 75 | 35 | 50 | 50 | 42 | 3 |
| 17.0 | 96 | | 35 | | 1 | 1 | 17 | 16 | 43 | 27 | 42 | 26 |
| 19.0 | 136 | 8 | 47 | | 10 | 2 | 3 | 20 | 22 | 10 | 34 | 52 |
| 21.0 | 105 | 33 | 87 | 2 | 57 | 1 | 0 | 5 | 16 | 1 | 17 | 22 |
| 23.0 | 39 | 150 | 44 | 22 | 182 | 12 | 10 | 34 | 11 | | 6 | 0 |
| 25.0 | 4 | 158 | 45 | 66 | 152 | 66 | 46 | 66 | 33 | 1 | | 0 |
| 27.0 | 5 | 41 | 84 | 62 | 67 | 202 | 219 | 123 | 82 | 1 | 1 | 1 |
| 29.0 | | 9 | 36 | 45 | 11 | 68 | 203 | 103 | 134 | | 0 | 0 |
| 31.0 | | 1 | 12 | 15 | 7 | 51 | 55 | 150 | 124 | | 1 | 2 |
| 33.0 | | | 1 | 3 | 1 | 11 | 3 | 95 | 95 | | 1 | |
| 35.0 | | | | 2 | 6 | 1 | 1 | 28 | 30 | | | |
| 37.0 | | | | 0 | 2 | 0 | | 11 | 9 | | | |
| 39.0 | | | | 1 | 1 | 3 | | 1 | | | | |

Publications of the Bay of Bengal Programme (BOBP)

The BOBP brings out six *types* of publications:

Reports (BOBP/REP/...) describe and analyze completed activities such as seminars, annual meetings of BOBP's Advisory Committee, and projects in member-countries for which BOBP inputs have ended

Working Papers (BOBP/WP/...) are progress reports that discuss the findings of ongoing BOBP work.

Manuals and Guides (BOBP/MAG/...) are instructional documents for specific audiences.

Miscellaneous Papers (BOBP/MIS/...) concern work not originated by BOBP staff or consultants – but which is relevant to the Programme's objectives.

Information Documents (BOBP/INF...) are bibliographies and descriptive documents on the fisheries of member-countries in the region.

Newsletters (*Ba, of Bengal News*), issued quarterly, contain illustrated articles and features in non-technical style on BOBP work and related subjects.

A list of publications follows.

Reports (BOBP/REP/...)

1. Report of the First Meeting of the Advisory Committee. Colombo, Sri Lanka, 28–29 October 1976. (Published as Appendix 1 of IOFC/DEV/78/44.1, FAO, Rome, 1978)
2. Report of the Second Meeting of the Advisory Committee. Madras, India, 29–30 June 1977. (Published as Appendix 2 of IOFC/DEV/78/44.1, FAO, Rome, 1978)
3. Report of the Third Meeting of the Advisory Committee. Chittagong, Bangladesh, 1–10 November 1978. Colombo, Sri Lanka, 1978. (Reissued Madras, India, September 1980)
4. Role of Women in Small-Scale Fisheries of the Bay of Bengal. Madras, India, October 1980.
5. Report of the Workshop on Social Feasibility in Small-Scale Fisheries Development. Madras, India, 3–8 September 1979. Madras, India, April 1980.
6. Report of the Workshop on Extension Service Requirements in Small-Scale Fisheries. Colombo, Sri Lanka, 8–12 October 1979. Madras, India, June 1980.
7. Report of the Fourth Meeting of the Advisory Committee. Phuket, Thailand, 27–30 November 1979. Madras, India, February 1980.
8. Pre-Feasibility Study of a Floating Fish Receiving and Distribution Unit for Dubla Char, Bangladesh. G. Eddie, M. T. Nathan. Madras, India, April 1980.
9. Report of the Training Course for Fish Marketing Personnel of Tamil Nadu. Madras, India, 3–14 December 1979. Madras, India, September 1980.
- 10.1 Report of the Consultation on Stock Assessment for Small-Scale Fisheries in the Bay of Bengal. Chittagong, Bangladesh, 16–21 June 1980. Volume 1: Proceedings. Madras, India, September 1980.
- 10.2 Report of the Consultation on Stock Assessment for Small-Scale Fisheries in the Bay of Bengal. Chittagong, Bangladesh, 16–21 June 1980. Volume 2: Papers. Madras, India, October 1980.
11. Report of the Fifth Meeting of the Advisory Committee. Penang, Malaysia, 4–7 November 1980. Madras, India, January 1981.
12. Report of the Training Course for Fish Marketing Personnel of Andhra Pradesh. Hyderabad, India, 11–26 November 1980. Madras, India, September 1981.
13. Report of the Sixth Meeting of the Advisory Committee. Colombo, Sri Lanka, 1–5 December 1981. Madras, India, February 1982.
14. Report of the First Phase of the "Aquaculture Demonstration for Small-Scale Fisheries Development Project" in Phang Nga Province, Thailand. Madras, India, March 1982.
15. Report of the Consultation-cum-Workshop on Development of Activities for Improvement of Coastal Fishing Families. Dacca, Bangladesh, October 27–November 6, 1981. Madras, India, May 1982.
16. Report of the Seventh Meeting of the Advisory Committee. New Delhi, India, January 17–21, 1983. Madras, India, March 1983.
17. Report of Investigations to Improve the Kattumaram of India's East Coast. Madras, India, July 1984.
18. Motorization of Country Craft, Bangladesh. Madras, India, July 1984.
19. Report of the Eighth Meeting of the Advisory Committee. Dhaka, Bangladesh, January 16–19, 1984. Madras, India, May 1984.

20. Coastal Aquaculture Project for Shrimp and Firtfish in Ban Merbok, Kedah, Malaysia. -Madras, India, December 1984.
21. Income-Earning Activities for Women from Fishing Communities in Sri Lanka. Edeltraud Drewes. Madras, India, September 1985.
22. Report of the Ninth Meeting of the Advisory Committee. Bangkok, Thailand, February 25—26, 1985. Madras, India, May 1985.
23. Summary Report of BOBP Fishing Trials and Demersal Resources Studies in Sri Lanka. Madras, India, March 1986.
24. Fisherwomen's Activities in Bangladesh: A Participatory Approach to Development. Patchanee Natpracha. Mkdiras, India, May 1986.
25. Attempts to Stimulate Development Activities in Fishing Communities of Adirampattinam, India. Madras, India, May 1986.
26. Report of the Tenth Meeting of the Advisory Committee. Male, Maldives 17—18 February 1986. Madras, India, April 1986.
27. Activating Fisherwomen for Development through Trained Link Workers in Tamil Nadu, India. Edeltraud Drewes. Madras, India, May 1986.
28. 8small-Scale Aquaculture Development Project in South Thailand: Results and Impact. e Drewes. Madras, India, May 1986.
29. Towards Shared Learning: An Approach to Non-formal Adult Education for Marine Fisherfolk of Tamil Nadu, India. L. S. Saraswathi and Patchanee Natpracha. Madras, India, July 1986.
30. Summary Report of Fishing Trials with Large-Mesh Driftnets in Bangladesh. Madras, India, May 1986.

Working Papers (BOBP/WP/....)

1. Investment Reduction and Increase in Service Life of Kattumaram Logs. R. Balan. Madras, India, February 1980.
2. Inventory of Kattumarams and their Fishing Gear in Andhra Pradesh and Tamil Nadu. t. R. Menon. Madras, India, October 1980.
3. Improvement of Large-Mesh Driftnets for Small-Scale Fisheries in Sri Lanka. C. Pajot. Madras, India, June 1980.
4. Inboard Motorisation of Small G.R.P. Boats in Sri Lanka. Madras, India, September 1980.
5. Improvement of Large-Mesh Driftnets for Small-Scale Fisheries in Bangladesh. C. Pajot. Madras, India, September 1980.
6. Fishing Trials with Bottom-Set Longlines in Sri Lanka. G. Pajot, K. T. Weerasooriya. Madras, India, September 1980.
7. Technical Trials of Beachcraft Prototypes in India. O. Gulbrandsen, G. P. Gowing, R. Ravikumar. Madras, India, October 1980.
8. Current Knowledge of Fisheries Resources in the Shelf Area of the Bay of Bengal. B. T. Antony Raja. Madras, India, September 1980.
9. Boatbuilding Materials for Small-Scale Fisheries in India. Madras, India, October 1980.
10. Fishing Trials with High-Opening Bottom Trawls in Tamil Nadu, India. C. Pajot, John Crockett. Madras, India, October 1980.
11. The Possibilities for Technical Cooperation between Developing Countries (TCDC) in Fisheries. E. H. Nichols. Madras, India, August 1981.
12. Trials in Bangladesh of Large-Mesh Driftnets of Light Construction. G. Pajot, T. K. Das. Madras, India, October 1981.
13. Trials of Two-Boat Bottom Trawling in Bangladesh. G. Pajot, J. Crockett. Madras, India, October 1982.
14. Three Fishing Villages in Tamil Nadu. Edeltraud Drewes. Madras, India, February 1982.
15. Effort Survey of Driftnet Fisheries in Bangladesh. M. Bergstrom. Madras, India, May 1982.
16. Further Trials with Bottom Longlines in Sri Lanka. Madras, India, July 1982.
17. Exploration of the Possibilities of Coastal Aquaculture Development in Andhra Pradesh. Soleh Sainsi, Sihar Siregar and Martono. Madras, India, September 1982.
18. Review of Brackishwater Aquaculture Development in Tamil Nadu. Kasemsant Chalayondeja and Avant Saraya. Madras, India, August 1982.
19. Coastal Village Development in Four Fishing Communities of Adirampattinam, Tamil Nadu, India. F. W. Blase. Madras, India, December 1982.
20. Further Trials of Mechanized Trawling for Food Fish in Tamil Nadu. O. Pajot, J. Crockett, S. Pandurangan, P. V. Ramamoorthy. Madras, India, December 1982.

21. Improved Deck Machinery and Layout for Small Coastal Trawlers. G. Pajot, **J. Crockett**, S. Pandurangan and **P. V. Ramamoorthy**. Madras, India, June 1983.
22. **The Impact of Management Training on the Performance of Marketing Officers in State Fisheries Corporations**, **U. Tietze**. Madras, India, June 1983.
23. Review of Experiences with and Present Knowledge about Fish Aggregating Devices. **M. Bergstrom**. Madras, India, November 1983.
24. Traditional Marine Fishing Craft and Gear of Orissa. **P. Mohapatra**. Madras, India, April 1986.
25. Fishing Craft Development in Kerala: Evaluation Report. **O. Gulbrandsen**. Madras, India, June 1984.
26. Commercial Evaluation of IND-13 Beachcraft at Uppada, India. R. Ravikumar. Madras, India, June 1964.
27. **Reducing Fuel Costs of Fishing Boats**. O. Gulbrandsen. Madras, India, July 1986.
28. Fishing Trials with Small-Mesh Driftnets in Bangladesh. G. Pajot and T. K. Das. Madras, India, March 1984.
29. Artisanal Marine Fisheries of Orissa: a Techno-Demographic Study. M. H. Kalavathy and U. Tietze. Madras, India, December 1984.
30. **Mackerels in the Malacca Straits**. Colombo, Sri Lanka, February 1985.
31. **Tuna Fishery in the EEZs of India, Maldives and Sri Lanka**. Colombo, Sri Lanka, February 1965.
32. **Pen Culture of Shrimp in the Backwaters of Killai, Tamil Nadu: A Study of Techno-economic and Social Feasibility**. Rathindra Nath Roy, Madras, India, January 1985.
33. **Factors that Influence the Role and Status of Fisherwomen**. Karuna Anbarasan. Madras, India, April 1985.
34. Pilot Survey of Set Bagnet Fisheries of Bangladesh. Abul Kashem. Madras, India, August 1985.
35. Pen Culture of Shrimp in the Backwaters of Killai, Tamil Nadu. M. Karim and S. Victor Chandra. Madras, India, May 1985.
36. Marine Fishery Resources of the Bay of Bengal. K. Sivasubramaniam. Colombo, Sri Lanka, October 1985.
37. **A Review of the Biology and Fisheries of *Hilsa ilisha* in the Upper Bay of Bengal**. B. T. Antony Raja. Colombo, Sri Lanka, October 1985.
38. **Credit for Fisherfolk: The Adirampattinam Experience**. R. S. Anbarasan and Ossie Fernandez. Madras, India, March 1986.
39. **The Organization of Fish Marketing in Madras Fishing Harbour**. M. H. Kalavathy. Madras, India, September 1985.
40. **Promotion of Bottom Set Longlining in Sri Lanka**. K. T. Weerasooriya, S. S. C. Pieris, M. Fonseka. Madras, India, August 1985.
41. **The Demersal Fisheries of Sri Lanka**. K. Sivasubramaniam and R. Maldeniya. Madras, India, December 1985.
42. **Fish Trap Trials in Sri Lanka**. (Based on a report by T. Hammerman). Madras, India, January 1986.
43. Demonstration of Simple Hatchery Technology for Prawns in Sri Lanka. Madras, India, June 1986.
44. **Pivoting Engine Installation for Beachlanding Boats**. A. Overa, R. Ravikumar. Madras, India, June 1986.
45. **Further Development of Beachlanding Craft in India and Sri Lanka**. A. Overa, R. Ravikumar, O. Gulbrandsen, G. Gowing. Madras, India, July 1986.
46. Experimental Shrimp Farming in Ponds in Polekurru, Andhra Pradesh, India. J. A. Janssen, T. Radhakrishna Murthy, B. V. Raghavulu, V. Sreekrishna. Madras, India, July 1986.
47. Growth and Mortality of the Malaysian Cockle (*Anaethara Granosa*) under Commercial Culture: Analysis through Length-Frequency Data. Ng Fong Oon. Madras, India, July 1986.

Manuals and Guides (BOBP/MAG/....)

1. Towards Shared Learning: Non-formal Adult Education for Marine Fisherfolk. Trainers' Manual. Madras, India, June 1985.
2. Towards Shared Learning: Non-formal Adult Education for Marine Fisherfolk. Animators' Guide. Madras, India, June 1985.
3. Fishery Statistics on the Microcomputer: A BASIC Version of Hasselblad's NORMSEP Program. D. Pauly, N. David, J. Hertel-Wulif. Madras, India, June 1986.

Miscellaneous Papers (BOBP/MIS/....)

1. Fishermen's Cooperatives in Kerala: A Critique. John Kurien. Madras, India, -October 1980.

2. Consultation on Social Feasibility of Coastal Aquaculture.
Madras, India, 26 November—1 December 1984. Madras, India, November 1985.

Information Documents (BOBP/INF/....)

1. Women and Rural Development in the Bay of Bengal Region: Information Sources.
Madras, India, February 1982.
2. Fish Aggregation Devices: Information Sources. Madras, India, February 1982.
3. Marine **Small-Scale Fisheries of India: A General Description. Madras, India, March 1983.**
4. Marine Small-Scale Fisheries of Andhra Pradesh: A General Description. Madras, India, June 1983.
5. Marine Small-Scale Fisheries of Tamil Nadu: A General Description. Madras, India, December 1983.
6. Marine **Small-Scale Fisheries of Sri Lanka: A General Description. Madras, India, November 1984.**
7. Marine Small-Scale Fisheries of Orissa: A General Description. Madras, India, December 1984.
8. Marine **Small-Scale Fisheries of Bangladesh: A General Description. Madras, India, September 1985.**
9. Food and Nutrition Status of Small-Scale Fishersfolk in India's East Coast States:
A Desk Review and Resource Investigation. V. Bhavani. Madras, India, April 1986.

Newsletters (Bay of Bengal News):

22 issues quarterly from January 1981 to June 1986.